

ASTRAZENECA PLC  
Form 6-K  
January 19, 2018

FORM 6-K

SECURITIES AND EXCHANGE COMMISSION  
Washington, D.C. 20549

Report of Foreign Issuer

Pursuant to Rule 13a-16 or 15d-16 of  
the Securities Exchange Act of 1934

For the month of January 2018

Commission File Number: 001-11960

AstraZeneca PLC

1 Francis Crick Avenue  
Cambridge Biomedical Campus  
Cambridge CB2 0AA  
United Kingdom

Indicate by check mark whether the registrant files or will file annual reports under cover of Form 20-F or Form 40-F.

Form 20-F  Form 40-F

Indicate by check mark if the registrant is submitting the Form 6-K in paper as permitted by Regulation S-T Rule 101(b)(1):

Indicate by check mark if the registrant is submitting the Form 6-K in paper as permitted by Regulation S-T Rule 101(b)(7): \_\_\_\_\_

Indicate by check mark whether the registrant by furnishing the information contained in this Form is also thereby furnishing the information to the Commission pursuant to Rule 12g3-2(b) under the Securities Exchange Act of 1934.

Yes  No

If "Yes" is marked, indicate below the file number assigned to the Registrant in connection with Rule 12g3-2(b):  
82- \_\_\_\_\_

AstraZeneca PLC

INDEX TO EXHIBITS

## ASTRAZENECA'S FASENRA RECEIVES APPROVAL IN JAPAN

19 January 2018 07:00 GMT

### FASENRA RECEIVES APPROVAL IN JAPAN

Approval based on Phase III WINDWARD programme that demonstrated significant reductions in asthma exacerbations, improvements in lung function and reductions in oral corticosteroid use from baseline, versus placebo

Fasenra is the first approved respiratory biologic with an 8-week maintenance dosing schedule

AstraZeneca and its global biologics research and development arm, MedImmune, today announced that the Japanese Ministry of Health, Labour and Welfare has approved Fasenra (benralizumab) as an add-on treatment for bronchial asthma in patients who continue to experience asthma exacerbations despite treatment with high-dose inhaled corticosteroid and other asthma controllers.

The approval is based on the results from the WINDWARD programme, including the pivotal Phase III exacerbation trials, SIROCCO and CALIMA, and the Phase III oral corticosteroid (OCS)-sparing trial, ZONDA.<sup>1</sup> Fasenra will be available as a fixed-dose subcutaneous injection in a prefilled syringe administered once every four weeks for the first three doses, and then once every eight weeks thereafter.<sup>1</sup>

Sean Bohan, Executive Vice President, Global Medicines Development and Chief Medical Officer at AstraZeneca, said: "The approval of Fasenra, our first respiratory biologic medicine, in Japan closely follows the recent US and EU decisions and brings us another step closer to achieving our ambition of transforming care for severe asthma patients around the world."

Fasenra binds directly to the IL-5a receptor on eosinophils, a type of white blood cell that are a normal part of the body's immune system,<sup>2</sup> and attracts natural killer cells to induce direct, rapid and near-complete depletion of eosinophils via apoptosis (programmed cell death).<sup>5,6,7</sup> Elevated levels of eosinophils, seen in about half of severe asthma patients, impact airway inflammation and airway hyper-responsiveness, resulting in increased asthma severity and symptoms, decreased lung function and increased risk of exacerbations.<sup>3,4</sup>

The Japanese approval follows US FDA approval in November 2017 and European Commission marketing authorisation in January 2018. Interactions with regulatory authorities in the rest of the world are on-going.

#### About Severe Asthma

Asthma affects approximately 315 million individuals worldwide. Up to 10% of asthma patients have severe asthma, which may be uncontrolled despite high doses of standard-of-care asthma controller medicines and can require the use of chronic OCS.<sup>3,9,10</sup>

Severe, uncontrolled asthma is debilitating and potentially fatal with patients experiencing frequent exacerbations and significant limitations on lung function and quality of life.<sup>11,12,13</sup> Severe, uncontrolled asthma has higher risk of mortality than severe asthma.<sup>12,14</sup>

Severe, uncontrolled asthma can lead to a dependence on OCS, with systemic steroid exposure potentially leading to serious short- and long-term adverse effects including weight gain, diabetes, osteoporosis, glaucoma, anxiety, depression, cardiovascular disease and immunosuppression.<sup>10,15,16,17,18</sup> There is also a significant physical and socio-economic burden of severe, uncontrolled asthma with these patients accounting for 50% of asthma-related costs.<sup>19</sup>

#### About Fasenera (benralizumab)

Fasenera is a monoclonal antibody that recruits natural killer cells to induce direct, rapid and near-complete depletion of eosinophils.<sup>7,20</sup> Depletion of circulating eosinophils is rapid, with an onset of action within 24 hours as confirmed in early Phase I/II trials.<sup>7,20,21</sup> Eosinophils are the biological effector cells in approximately 50% of asthma patients, leading to frequent exacerbations, impaired lung function and asthma symptoms.<sup>3,4</sup>

Fasenera is now approved in the US, EU and Japan, and under regulatory review in several other jurisdictions.

Fasenera is the foundation of AstraZeneca's respiratory biologics portfolio of potential new medicines targeting underlying causes of respiratory disease. Fasenera is also being evaluated in chronic obstructive pulmonary disease (COPD) with data readout expected in the second half of 2018.<sup>22</sup>

Fasenera was developed by AstraZeneca with MedImmune, the company's global biologics research and development arm and is in-licensed from BioWa, Inc., a wholly-owned subsidiary of Kyowa Hakko Kirin Co., Ltd., Japan.

#### About the WINDWARD Programme

The WINDWARD programme in asthma is made up six Phase III trials, including SIROCCO, CALIMA, ZONDA, BISE, BORA and GREGALE.<sup>22</sup> The two pivotal trials SIROCCO and CALIMA, are randomised, double-blinded, parallel-group, placebo-controlled trials designed to evaluate the efficacy and safety of a regular, subcutaneous administration of Fasenera (fixed 30mg dose) for up to 56-weeks in exacerbation-prone adult and adolescent patients 12 years of age and older.<sup>5,6</sup>

A total of 2,510 patients (1,204 in SIROCCO and 1,306 in CALIMA) received standard-of-care medicines (including high-dosage inhaled corticosteroids and long-acting beta2-agonists) and were randomised globally to receive either Fasenera 30mg every 4-weeks; Fasenera 30mg every 4-weeks for the first three doses followed by 30mg every 8-weeks; or placebo administered via subcutaneous injection using an accessorised pre-filled syringe.<sup>5,6,23</sup>

In SIROCCO and CALIMA, patients with severe, uncontrolled eosinophilic asthma receiving Fasenera experienced a significant reduction in asthma exacerbations and improvement in their lung function and asthma symptoms compared to patients receiving placebo, on top of their standard medicines.<sup>5,6</sup> The most commonly reported adverse reactions during treatment were headache (8%) and pharyngitis (3%). Other common adverse reactions included fever, hypersensitivity reactions and injection site reactions.<sup>1,5,6</sup>

A pooled post-hoc analysis of the SIROCCO and CALIMA trials, demonstrated an association between enhanced Fasenera efficacy and certain easily identifiable clinical features of severe eosinophilic asthma, including baseline blood eosinophil counts, history of more frequent exacerbations, chronic OCS use and a history of nasal polyposis.<sup>23</sup>

The third registrational trial, ZONDA, demonstrated a statistically-significant and clinically-meaningful reduction in daily-maintenance OCS use compared with placebo for patients with severe, uncontrolled OCS-dependent eosinophilic asthma receiving Fasenera. The results were published in the New England Journal of Medicine in May 2017.<sup>24</sup>

In addition to WINDWARD, the Phase III VOYAGER programme is currently underway, which is evaluating the efficacy and safety of Fasenera in patients with severe COPD.<sup>22</sup>

#### About AstraZeneca in Respiratory Disease

Respiratory disease is one of AstraZeneca's main therapy areas, and the Company has a growing portfolio of medicines that reached more than 18 million patients in 2017. AstraZeneca's aim is to transform asthma and COPD treatment through inhaled combinations at the core of care, biologics for the unmet needs of specific patient populations, and scientific advancements in disease modification.

The Company is building on a 40-year heritage in respiratory disease and AstraZeneca's capability in inhalation technology spans pressurised metered-dose inhalers and dry powder inhalers, as well as the Aerosphere Delivery Technology. The company's biologics include Fasenra, (anti-eosinophil, anti-IL-5r ), which is now approved in the US, EU, Japan, and is under regulatory review in other jurisdictions, and tezepelumab (anti-TSLP), which successfully achieved its Phase IIb primary and secondary endpoints and has initiated Phase III. AstraZeneca's research is focused on addressing underlying disease drivers focusing on the lung epithelium, lung immunity and lung regeneration.

#### About MedImmune

MedImmune is the global biologics research and development arm of AstraZeneca, a global, innovation-driven biopharmaceutical business that focuses on the discovery, development and commercialisation of small molecule and biologic prescription medicines. MedImmune is pioneering innovative research and exploring novel pathways across Oncology, Respiratory, Cardiovascular & Metabolic Diseases, and Infection and Vaccines. The MedImmune headquarters is located in Gaithersburg, Md., one of AstraZeneca's three global R&D centres, with additional sites in Cambridge, UK and Mountain View, Calif. For more information, please visit [www.medimmune.com](http://www.medimmune.com)

#### About AstraZeneca

AstraZeneca is a global, science-led biopharmaceutical company that focuses on the discovery, development and commercialisation of prescription medicines, primarily for the treatment of diseases in three therapy areas - Oncology, Cardiovascular & Metabolic Diseases and Respiratory. The Company also is selectively active in the areas of autoimmunity, neuroscience and infection. AstraZeneca operates in over 100 countries and its innovative medicines are used by millions of patients worldwide.

For more information, please visit [www.astrazeneca.com](http://www.astrazeneca.com) and follow us on Twitter @AstraZeneca.

#### Media Relations

Esra Erkal-Paler	UK/Global	+44 203 749 5638
Karen Birmingham	UK/Global	+44 203 749 5634
Rob Skelding	UK/Global	+44 203 749 5821
Matt Kent	UK/Global	+44 203 749 5906
Gonzalo Viña	UK/Global	+44 203 749 5916
Jacob Lund	Sweden	+46 8 553 260 20
Michele Meixell	US	+1 302 885 2677

#### Investor Relations

Thomas Kudsk Larsen		+44 203 749 5712
Craig Marks	Finance, Fixed Income, M&A	+44 7881 615 764
Henry Wheeler	Oncology	+44 203 749 5797
Mitchell Chan	Oncology; Other	+1 240 477 3771

## Edgar Filing: ASTRAZENECA PLC - Form 6-K

Christer Gruvris	Brilinta; Diabetes	+44 203 749 5711
Nick Stone	Respiratory; Renal	+44 203 749 5716
US toll free		+1 866 381 7277

Adrian Kemp  
Company Secretary  
AstraZeneca PLC

### References

1. Fasenna (benralizumab) Package Insert. AstraZeneca PLC, 2018.
2. Mukherjee M, Sehmi R, Nair P. Anti-IL5 therapy for asthma and beyond. *World Allergy Organ J.* 2014;7:32.
3. Wenzel S. Severe asthma in adults. *Am J Respir Crit Care Med.* 2005;172:149-160.
4. Zhang, JY and Wenzel, SE. Tissue and BAL based biomarkers in asthma. *Immunol Allergy Clin North Am.* 2007; 27: 623-632 (vi.).
5. FitzGerald MJ, Bleecker E, Parameswaran N, et al. Benralizumab, an anti-interleukin-5 receptor a monoclonal antibody, as add-on treatment for patients with severe, uncontrolled, eosinophilic asthma (CALIMA): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet.* 2016. Volume 388, Issue 10056, 2128 - 2141.
6. Bleecker ER, FitzGerald MJ, Chanez P, et al. Efficacy and safety of benralizumab for patients with severe asthma uncontrolled with high-dosage inhaled corticosteroids and long-acting b2-agonists (SIROCCO): a randomised, multicentre, placebo-controlled phase 3 trial. *Lancet.* 2016. Volume 388, Issue 10056, 2115 - 2127.
7. Kolbeck R, Kozhich A, Koike M, et al. MEDI-563, a humanized anti-IL-5 receptor a mAb with enhanced antibody-dependent cell-mediated cytotoxicity function. *J Allergy Clin Immunol.* 2010 Jun;125(6):1344-1353.e2.
8. Fasenna Prescribing Information. AstraZeneca Pharmaceuticals LP.
9. To T, Stanojevic S, Moores G, et al. Global asthma prevalence in adults: findings from cross-sectional world health survey. *BioMed Central Public Health.* 2012: 12(204).
10. Global Initiative for Asthma (GINA). Online appendix. Global strategy for asthma management and prevention. Updated 2017. Available from: <http://ginasthma.org/2017-gina-report-global-strategy-for-asthma-management-and-prevention/>. Last accessed November 2017.
11. Price D, Fletcher M, van der Molen T. Asthma control and management in 8,000 European patients: the REcognise Asthma and LInk to Symptoms and Experience (REALISE) survey. *NPJ Prim Care Respir Med* 2014; 12; 24: 14009.
12. Peters SP, Ferguson G, Deniz Y, et al. Uncontrolled asthma: a review of the prevalence, disease burden and options for treatment. *Respir Med.* 2006: 100(7):1139-51.
13. Adelphi Real World Respiratory Disease Specific Programme. 2012-2014. [Asthma patient data file], Bollington, UK. Unpublished raw data, cited with permission.
14. Fernandes AG, Souza-Machado C, Coelho RC et al. Risk factors for death in patients with severe asthma. *J Bras Pneumol.* 2014; 40(4): 364-372.
15. Hyland ME, Whalley B, Jones RC, Masoli M. A qualitative study of the impact of severe asthma and its treatment showing that treatment burden is neglected in existing asthma assessment scales. *Qual Life Res.* 2015 Mar;24(3):631-9. doi: 10.1007/s11136-014-0801-x. Epub 2014 Sep 9.
16. Global Initiative for Asthma (GINA). Online appendix. Global strategy for asthma management and prevention. Updated 2017. Available from: <http://ginasthma.org/2017-gina-report-global-strategy-for-asthma-management-and-prevention/>. Last accessed December 2017.
17. Iribarren C, Tolstykh IV, Miller MK, et al. Adult asthma and risk of coronary heart disease, cerebrovascular disease, and heart failure: a prospective study of 2 matched cohorts. *Am J Epidemiol.* 2012;176:1014-1024.

Edgar Filing: ASTRAZENECA PLC - Form 6-K

18. Zazzali JL, Broder MS, Omachi TA, Chang E, Sun GH, Raimundo K. Risk of corticosteroid-related adverse events in asthma patients with high oral corticosteroid use. *Allergy Asthma Proc.* 2015 Jul-Aug;36(4):268-74. doi: 10.2500/aap.2015.36.3863.
19. World Allergy Organization (WAO). The management of severe asthma: economic analysis of the cost of treatments for severe asthma. Available from: [http://www.worldallergy.org/educational\\_programs/world\\_allergy\\_forum/anaheim2005/blaiss.php](http://www.worldallergy.org/educational_programs/world_allergy_forum/anaheim2005/blaiss.php). Last accessed May 2017.
20. Pham TH, Damera G, Newbold P, Ranade K. Reductions in eosinophil biomarkers by benralizumab in patients with asthma. *Respir Med.* 2016; 111:21-29.
21. Laviolette M, Gossage DL, Gauvreau G, et al. Effects of benralizumab on airway eosinophils in asthmatic patients with sputum eosinophilia. *J Allergy Clin Immunol.* 2013 Nov; 132(5): 1086 - 1096.e5.
22. AstraZeneca. Clinical Trials Appendix Q2 2017 Results Update. Available at [https://www.astrazeneca.com/content/dam/az/PDF/2017/H1/H1\\_2017\\_Results\\_Clinical\\_Trial\\_Appendix.pdf](https://www.astrazeneca.com/content/dam/az/PDF/2017/H1/H1_2017_Results_Clinical_Trial_Appendix.pdf). Last Accessed November 2017.
23. FitzGerald JM, Bleecker ER, Menzies-Gow A, Zangrilli JG, Hirsch I, Metcalfe P, Newbold P, Goldman M. Predictors of enhanced response with benralizumab for patients with severe asthma: pooled analysis of the SIROCCO and CALIMA studies. *Lancet Respir Med.* 2017 Sep 11.
24. Nair P, Wenzel S, Rabe KF, Bourdin A, Lugogo NL, et al. Oral glucocorticoid-sparing effect of benralizumab in severe asthma. *N Engl J Med.* 2017; 376:2448-58.

SIGNATURES

Pursuant to the requirements of the Securities Exchange Act of 1934, the Registrant has duly caused this report to be signed on its behalf by the undersigned, thereunto duly authorized.

AstraZeneca PLC

Date: 19 January 2018

By: /s/ Adrian Kemp

Name: Adrian Kemp

Title: Company Secretary